

Correlated scattering of photons from a trapped Bose-Einstein condensate

Hiroki Saito and Masahito Ueda

*Department of Physical Electronics, Hiroshima University, Higashi-Hiroshima 739-8527, Japan
and CREST, Japan Science and Technology Corporation (JST), Higashi-Hiroshima 739-8527, Japan*

The second-order coherence of photons scattered from a trapped Bose-Einstein condensate is found to be enhanced for the scattering angles that are either the same or symmetrical with respect to the direction of laser propagation. The enhancement occurs periodically with respect to the time interval between photons, and becomes less and less pronounced with increasing the strength of the interaction between atoms.

03.75.Fi, 42.50.Vk, 42.50.Gy, 05.30.Jp

I. INTRODUCTION

Realization of Bose-Einstein condensation (BEC) in trapped atomic gases [1–4] has created an interdisciplinary subfield of physics called coherent matter wave. Developments of optical imaging techniques such as absorption imaging and a nondestructive *in situ* method [5] have enabled us to observe interference between matter waves [6] and dynamics of BEC [7,8]. In a remarkable recent experiment, Hau *et al.* [9] demonstrated that it is possible to make photons propagate through BEC at a speed of 17 m/s, which suggests that the index of refraction of BEC is 14 orders of magnitude greater than that of a glass fiber. In the present paper we predict yet another unique feature of BEC concerning correlated photon scattering.

Various aspects of the optical response of BEC have been predicted [10–20]: a broad resonance [13,15], an extra structure in the spectrum [16,19], response to intense short pulses [12,17], non-Lorentzian line shapes [15,20], and a refractive index of a homogeneous gas [18]. In the above work, kinetic motion of atoms is either ignored or not explicitly taken into account because its time scale is usually much slower than that of optical processes. However, optical signatures of BEC are expected to appear also in the time scale of kinetic motion, since the characteristic feature of BEC is the degeneracy of the kinetic degrees of freedom as well as the atomic internal degree of freedom.

The aim of the present paper is to study correlations of photons scattered from trapped BEC that is irradiated by a weak far-off-resonant laser field, and to show that they exhibit some characteristic features associated with kinetic motion of the atoms in the trap. By scattering a photon, the condensate suffers a recoil, which affects the scattering of the subsequent photons. As a consequence a correlation arises between a pair of photons with respect to their scattering angles and time intervals. We

show that the presence of the condensate enhances the probability of a pair of photons to be scattered into either the same direction or symmetrical directions about laser propagation, and the enhancement occurs periodically with respect to the time interval, where the period is set by the frequency of the trap.

The interaction between atoms cannot be ignored, for the profile of the condensate is significantly altered even for a few hundred atoms [21,22]. While in Ref. [20] effects of the atom-atom interaction on the photon scattering are taken into account only through the atomic density profile, the dynamical effect of the interaction on the optical response has not been discussed. We take into account the atom-atom interaction using the Bogoliubov approximation [23], and show that the correlations between the scattered photons are deteriorated by the interaction.

This paper is organized as follows. Section II formulates the problem and derives an expression of the electric-field operator far from BEC irradiated by a laser field. Section III studies coherence properties of scattered photons. Section IV discusses effects of the atom-atom interaction on the scattering of photons, and Sec. V concludes this paper.

II. FORMULATION OF THE PROBLEM

We consider a situation in which trapped atoms are irradiated by a laser field which is sufficiently detuned from the resonant frequency of the atoms so that each photon is not scattered by the atoms more than once. Photons are scattered from the atoms and detected by photon counters. Quantum-statistical properties of the scattered photons can be studied through the electric-field operator $\hat{\mathbf{E}}(\mathbf{r}, t)$ in the Heisenberg representation. We derive here an expression of $\hat{\mathbf{E}}(\mathbf{r}, t)$ at the position far away from the trapped atoms compared with the spread of their wave functions.

The Hamiltonian for the photon field is given by

$$\hat{H}_f = \sum_{\mathbf{k}\sigma} \hbar\omega_{\mathbf{k}} \hat{a}_{\mathbf{k}\sigma}^\dagger \hat{a}_{\mathbf{k}\sigma}, \quad (1)$$

where $\hat{a}_{\mathbf{k}\sigma}^\dagger$ and $\hat{a}_{\mathbf{k}\sigma}$ are the creation and annihilation operators of a photon with wave vector $\mathbf{k}$ and polarization σ . The electric-field operator is expanded in terms of the creation and annihilation operators as

$$\hat{\mathbf{E}}(\mathbf{r}) = i \sum_{\mathbf{k}\sigma} \sqrt{\frac{\hbar\omega_{\mathbf{k}}}{2\varepsilon_0 V}} \left(\boldsymbol{\epsilon}_{\mathbf{k}\sigma} e^{i\mathbf{k}\cdot\mathbf{r}} \hat{a}_{\mathbf{k}\sigma} - \boldsymbol{\epsilon}_{\mathbf{k}\sigma}^* e^{-i\mathbf{k}\cdot\mathbf{r}} \hat{a}_{\mathbf{k}\sigma}^\dagger \right), \quad (2)$$

where $\epsilon_{\mathbf{k}\sigma}$ is the polarization vector of photons. The internal states of two-level atoms are described by the Hamiltonian

$$\hat{H}_{\text{ai}} = \sum_{j=1}^N \frac{\hbar\omega_A}{2} \hat{\sigma}_{jz}, \quad (3)$$

where N is the number of atoms, $\hbar\omega_A$ is the transition energy of the internal levels, and the operator $\hat{\sigma}_{jz}$ is the Pauli operator which takes on the eigenvalue 1 (or -1) when the j th atom is in the excited (or ground) internal level. The Hamiltonian corresponding to the kinetic degrees of freedom of the atoms is given by

$$\hat{H}_{\text{ak}} = \sum_{j=1}^N \left[-\frac{\hbar^2}{2m} \nabla_j^2 + V_t(\mathbf{R}_j) \right] + \sum_{j \neq j'}^N \frac{2\pi\hbar^2 a}{m} \delta(\mathbf{R}_j - \mathbf{R}_{j'}), \quad (4)$$

where m is the mass of the atom and $V_t(\mathbf{R})$ is the trap potential. The last term in Eq. (4) describes the Fermi's contact type of interaction between hard-core atoms, which is characterized by the s-wave scattering length a .

We assume the electric-dipole interaction between the atoms and the photon field which is described by the Hamiltonian

$$\hat{H}_{\text{ed}} = - \sum_{j=1}^N \hat{\mathbf{D}}_j \cdot \hat{\mathbf{E}}(\mathbf{R}_j). \quad (5)$$

The dipole operator denoted as $\hat{\mathbf{D}}_j$ has the form

$$\hat{\mathbf{D}}_j \equiv \mathbf{d} \hat{\sigma}_{j+} + \mathbf{d}^* \hat{\sigma}_{j-}, \quad (6)$$

where $\mathbf{d}$ is the electric-dipole matrix element, and $\hat{\sigma}_{j+}$ and $\hat{\sigma}_{j-}$ are the raising and lowering operators of the internal state of the j th atom.

The Hamiltonian for the atom-laser interaction is given by

$$\hat{H}_L = \sum_{j=1}^N \left[\mathcal{E}_L(\epsilon_L \cdot \mathbf{d}) e^{-i(\omega_L t - \mathbf{k}_L \cdot \mathbf{R}_j)} \hat{\sigma}_{j+} + \mathcal{E}_L^*(\epsilon_L \cdot \mathbf{d})^* e^{i(\omega_L t - \mathbf{k}_L \cdot \mathbf{R}_j)} \hat{\sigma}_{j-} \right], \quad (7)$$

where $\mathcal{E}_L$, ϵ_L , ω_L , and $\mathbf{k}_L$ denote the amplitude, the polarization vector, the frequency, and the wave vector of the laser field, respectively. Since the optical processes of the internal states of the atoms are much faster than those of the kinetic motion, we may adiabatically eliminate the dynamics of the internal state. When the detuning $\delta \equiv \omega_L - \omega_A$ is much larger than both the atom-laser field coupling and the collective linewidth Γ of the atoms [13,15], i.e.,

$$|\delta| \gg \frac{|\mathcal{E}_L \mathbf{d}|}{\hbar}, \quad |\delta| \gg \Gamma, \quad (8)$$

the probability that the atoms are in the excited state $|e\rangle$ is much smaller than that in the ground state $|g\rangle$. In this case, the atomic internal state may be assumed to be in the ground state of the Hamiltonian $\hat{H}_{\text{ai}} + \hat{H}_L$, which is given, up to the first order in $|\mathcal{E}_L \mathbf{d}|/\hbar\delta$, by

$$\prod_{j=1}^N \left[e^{i(\omega_L t - \mathbf{k}_L \cdot \mathbf{R}_j)/2} |g\rangle_j + \frac{|\mathcal{E}_L \mathbf{d}|}{\hbar\delta} e^{-i(\omega_L t - \mathbf{k}_L \cdot \mathbf{R}_j)/2} |e\rangle_j \right]. \quad (9)$$

Taking the expectation value of $\hat{H}_{\text{ed}}$ in Eq. (5) with respect to the atomic internal state (9), we obtain the effective Hamiltonian describing the interaction between photons and kinetic motion of the atoms via the atomic internal levels as

$$\hat{H}'_{\text{ed}} = \sum_{j=1}^N \frac{|\mathcal{E}_L \mathbf{d}|}{\hbar\delta} \left[e^{i(\omega_L t - \mathbf{k}_L \cdot \mathbf{R}_j)} \mathbf{d} + e^{-i(\omega_L t - \mathbf{k}_L \cdot \mathbf{R}_j)} \mathbf{d}^* \right] \cdot \hat{\mathbf{E}}(\mathbf{R}_j). \quad (10)$$

In the following discussions we will employ the effective Hamiltonian $\hat{H}_{\text{eff}} = \hat{H}_f + \hat{H}_{\text{ak}} + \hat{H}'_{\text{ed}}$ to describe the dynamics of photons and kinetic motion of the atoms.

The equation of motion for the electric-field operator in the Heisenberg representation is obtained from the Hamiltonian $\hat{H}_{\text{eff}}$ as

$$\left(\nabla^2 - \frac{\partial^2}{c^2 \partial t^2} \right) \hat{\mathbf{E}}(\mathbf{r}, t) = \frac{i}{\hbar} [\hat{H}'_{\text{ed}}(t), \nabla^2 \hat{\mathbf{A}}(\mathbf{r}, t)], \quad (11)$$

where the vector potential is given by

$$\hat{\mathbf{A}}(\mathbf{r}, t) = \sum_{\mathbf{k}\sigma} \sqrt{\frac{\hbar}{2\varepsilon_0 \omega_k V}} \left[e^{i\mathbf{k} \cdot \mathbf{r}} \epsilon_{\mathbf{k}\sigma} \hat{a}_{\mathbf{k}\sigma}(t) + e^{-i\mathbf{k} \cdot \mathbf{r}} \epsilon_{\mathbf{k}\sigma}^* \hat{a}_{\mathbf{k}\sigma}^\dagger(t) \right]. \quad (12)$$

The equation of motion (11) can be solved, giving

$$\hat{\mathbf{E}}(\mathbf{r}, t) = \hat{\mathbf{E}}_{\text{vac}}(\mathbf{r}, t) + \hat{\mathbf{E}}_{\text{s}}(\mathbf{r}, t), \quad (13)$$

where $\hat{\mathbf{E}}_{\text{vac}}(\mathbf{r}, t) = e^{\frac{i}{\hbar} \hat{H}_f t} \hat{\mathbf{E}}(\mathbf{r}) e^{-\frac{i}{\hbar} \hat{H}_f t}$ is the vacuum field and

$$\hat{\mathbf{E}}_{\text{s}}(\mathbf{r}, t) = -\frac{i}{4\pi\hbar} \int d\mathbf{r}' dt' \frac{[\hat{H}'_{\text{ed}}(t'), \nabla^2 \hat{\mathbf{A}}(\mathbf{r}', t')]}{|\mathbf{r} - \mathbf{r}'|} \times \delta(t - t' - |\mathbf{r} - \mathbf{r}'|/c) \quad (14)$$

describes the source field scattered from the atoms. Evaluating the commutator and carrying out the integral with respect to t' , the positive frequency part of the source field (14) becomes

$$\begin{aligned} \hat{\mathbf{E}}_{\text{s}}^{(+)}(\mathbf{r}, t) &= -\frac{1}{4\pi\varepsilon_0 V} \frac{|\mathcal{E}_L \mathbf{d}|}{\hbar\delta} \int d\mathbf{r}' \frac{e^{-i\omega_L t_r}}{|\mathbf{r} - \mathbf{r}'|} \\ &\times \sum_{\mathbf{k}} \mathbf{k} \times (\mathbf{d} \times \mathbf{k}) e^{-i\mathbf{k} \cdot \mathbf{r}'} \sum_{j=1}^N e^{i(\mathbf{k}_L + \mathbf{k}) \cdot \mathbf{R}_j(t_r)}, \end{aligned} \quad (15)$$

where $t_r \equiv t - |\mathbf{r} - \mathbf{r}'|/c$ and $\mathbf{R}_j(t_r) \equiv e^{\frac{i}{\hbar}\hat{H}_{\text{eff}}t_r}\mathbf{R}_je^{-\frac{i}{\hbar}\hat{H}_{\text{eff}}t_r}$. Since $|\mathcal{E}_L\mathbf{d}|/\hbar\delta \ll 1$, it is sufficient to keep only the terms first order in $|\mathcal{E}_L\mathbf{d}|/\hbar\delta$, which amounts to disregarding the multiple scattering and the dipole-dipole interaction. Consequently, we can make an approximation

$$\mathbf{R}_j(t_r) \simeq e^{\frac{i}{\hbar}\hat{H}_{\text{ak}}t}\mathbf{R}_je^{-\frac{i}{\hbar}\hat{H}_{\text{ak}}t} \equiv \mathbf{R}_j(t), \quad (16)$$

where we ignore the time $|\mathbf{r} - \mathbf{r}'|/c$ it takes photons to propagate from the atoms to the detector as it is much smaller than the time scale of the kinetic motion of the atoms. The integrand of the source field (15) contributes only when $\mathbf{r}'$ falls within the spread of the atomic wave function. Therefore, if we choose the origin of the coordinate system at the center of the trap, the distance $|\mathbf{r}|$ from the atom to the detector is much larger than $|\mathbf{r}'|$, and hence $t_r \simeq t - (r/c)(1 - \mathbf{r} \cdot \mathbf{r}'/r^2)$ and $1/|\mathbf{r} - \mathbf{r}'| \simeq 1/r$. We thus obtain

$$\begin{aligned} \hat{\mathbf{E}}_s^{(+)}(\mathbf{r}, t) &= -\frac{1}{4\pi\epsilon_0 r V} \frac{|\mathcal{E}_L\mathbf{d}|}{\delta\hbar} e^{-i\omega_L t} \int d\mathbf{r}' \sum_{\mathbf{k}} \mathbf{k} \times (\mathbf{d} \times \mathbf{k}) \\ &\quad \times e^{ikr} e^{i(k_L\tilde{\mathbf{r}} - \mathbf{k}_L) \cdot \mathbf{r}'} \sum_{j=1}^N e^{i(\mathbf{k}_L + \mathbf{k}) \cdot \mathbf{R}_j(t)} \\ &= -\frac{k_L^2}{4\pi\epsilon_0 r} \frac{|\mathcal{E}_L\mathbf{d}|}{\delta\hbar} \mathbf{n} \times (\mathbf{d} \times \mathbf{n}) e^{-i\omega_L t + ik_L r} \\ &\quad \times \sum_{j=1}^N e^{i(\mathbf{k}_L - k_L\mathbf{n}) \cdot \mathbf{R}_j(t)} \\ &\equiv \mathbf{F}(\mathbf{r}, t) \sum_{j=1}^N e^{i\Delta\mathbf{k} \cdot \mathbf{R}_j(t)}, \end{aligned} \quad (17)$$

where $\mathbf{n} \equiv \mathbf{r}/|\mathbf{r}|$ and $\Delta\mathbf{k} \equiv \mathbf{k}_L - k_L\mathbf{n}$. The expression of the source-field operator (17) consists of the factor $\mathbf{F}(\mathbf{r}, t)$ which depends on the spatial configuration of the system and the operator that transfers a recoil momentum $\hbar\Delta\mathbf{k}$ to the atoms.

III. FIRST AND SECOND-ORDER COHERENCE OF PHOTONS SCATTERED FROM A NONINTERACTING CONDENSATE

In this section we consider an ideal Bose gas, where the atom-atom interaction is absent. We write the eigenfunctions of the Hamiltonian $\hat{H}_{\text{ak}}$ as $f_n(\mathbf{R})$, where n is an index for the eigenstates with $n = 0$ referring to the ground state.

It is convenient to discuss the problem in the second-quantized formalism. We expand the atomic field operator as

$$\hat{\psi}(\mathbf{R}) = \sum_n f_n(\mathbf{R}) \hat{b}_n, \quad (18)$$

where $\hat{b}_n$ annihilates an atom in the n th eigenstate, and satisfies the Bose commutation relation $[\hat{b}_n, \hat{b}_{n'}^\dagger] = \delta_{nn'}$. The Hamiltonian $\hat{H}_{\text{ak}}$ can then be rewritten as

$$\hat{H}_{\text{ak}} = \sum_n \hbar\omega_n \hat{b}_n^\dagger \hat{b}_n, \quad (19)$$

where $\hbar\omega_n$ is the n th eigenenergy. The second-quantized form of the operator $\sum_{j=1}^N e^{i\Delta\mathbf{k} \cdot \mathbf{R}_j(t)}$ in Eq. (17) is given by

$$\begin{aligned} \hat{\mathcal{D}}(\Delta\mathbf{k}, t) &\equiv \int d\mathbf{R} \hat{\psi}^\dagger(\mathbf{R}, t) e^{i\Delta\mathbf{k} \cdot \mathbf{R}} \hat{\psi}(\mathbf{R}, t) \\ &= \sum_{nn'} \langle n' | e^{i\Delta\mathbf{k} \cdot \mathbf{R}} | n \rangle \hat{b}_n^\dagger \hat{b}_n e^{i(\omega_{n'} - \omega_n)t}, \end{aligned} \quad (20)$$

where $\hat{\psi}(\mathbf{r}, t) \equiv e^{\frac{i}{\hbar}\hat{H}_{\text{ak}}t} \hat{\psi}(\mathbf{R}) e^{-\frac{i}{\hbar}\hat{H}_{\text{ak}}t}$, and the bracket $\langle n' | e^{i\Delta\mathbf{k} \cdot \mathbf{R}} | n \rangle$ is a shorthand notation of $\int d\mathbf{R} f_{n'}^*(\mathbf{R}) e^{i\Delta\mathbf{k} \cdot \mathbf{R}} f_n(\mathbf{R})$. The source-field operator (17) is expressed as $\hat{\mathbf{E}}^{(+)}(\mathbf{r}, t) = \mathbf{F}(\mathbf{r}, t) \hat{\mathcal{D}}(\Delta\mathbf{k}, t)$.

At zero temperature all N atoms are Bose-Einstein condensed in the ground-state level of the trap potential. We assume an isotropic harmonic trap $V_t(\mathbf{R}) = m\omega^2 R^2/2$ for simplicity. The expectation value of the electric field at the position $\mathbf{r}$ becomes

$$\begin{aligned} \langle \hat{\mathbf{E}}^{(+)}(\mathbf{r}, t) \rangle &= \mathbf{F}(\mathbf{r}, t) N \langle 0 | e^{i\Delta\mathbf{k} \cdot \mathbf{R}} | 0 \rangle \\ &= \mathbf{F}(\mathbf{r}, t) N e^{-|d\Delta\mathbf{k}|^2/4}, \end{aligned} \quad (21)$$

where $d \equiv (\hbar/m\omega)^{1/2}$ denotes the characteristic length scale of the trap. The exponential factor in Eq. (21) is a decreasing function of the scattering angle.

Suppose that we detect the number of scattered photons in a configuration schematically illustrated in Fig. 1. The position of the i th detector is denoted as $\mathbf{r}_i$ and the scattering angle as measured from the direction of laser propagation is denoted as θ_i . The number of photons counted by the detector is proportional to the normally-ordered intensity of the field [24]

$$I(\mathbf{r}, t) \equiv \langle \hat{\mathbf{E}}^{(-)}(\mathbf{r}, t) \cdot \hat{\mathbf{E}}^{(+)}(\mathbf{r}, t) \rangle. \quad (22)$$

Substituting Eq. (17) into this gives

$$I(\mathbf{r}, t) = |\mathbf{F}(\mathbf{r}, t)|^2 N \left[1 + (N-1) e^{-|d\Delta\mathbf{k}|^2/2} \right]. \quad (23)$$

The first term on the right-hand side is proportional to the number of atoms N and the angular dependence lies solely in $|\mathbf{F}(\mathbf{r}, t)|^2$. The second term is proportional to $N(N-1)$ and contains a forward-scattering factor of $e^{-|d\Delta\mathbf{k}|^2/2}$. In fact, in experiments of Ref. [5], a forward scattering that satisfies $|d\Delta\mathbf{k}| \lesssim 1$ was observed at the onset of BEC. The degree of first-order coherence is defined by [24]

$$g^{(1)}(\mathbf{r}_1, t_1; \mathbf{r}_2, t_2) \equiv \frac{\langle \hat{\mathbf{E}}^{(-)}(\mathbf{r}_1, t_1) \cdot \hat{\mathbf{E}}^{(+)}(\mathbf{r}_2, t_2) \rangle}{\sqrt{I(\mathbf{r}_1, t_1) I(\mathbf{r}_2, t_2)}}. \quad (24)$$




FIG. 1. Schematic of a scattering experiment. The trapped Bose-Einstein condensate (BEC) is irradiated by a weak and far off-resonant laser. The scattered photons are detected by the detectors set around the BEC. The positions of the detectors are denoted as $\mathbf{r}_i$ and the scattering angles as θ_i ($i = 1, 2$).

Substituting Eq. (17) into this gives

$$|g^{(1)}(\mathbf{r}_1, t_1; \mathbf{r}_2, t_2)| = \frac{e^{-|d\Delta\mathbf{k}_1|^2/4 - |d\Delta\mathbf{k}_2|^2/4} |e^{d^2\Delta\mathbf{k}_1 \cdot \Delta\mathbf{k}_2 e^{-i\omega(t_1-t_2)/2} + N - 1}|}{[1 + (N-1)e^{-|d\Delta\mathbf{k}_1|^2/2}]^{\frac{1}{2}} [1 + (N-1)e^{-|d\Delta\mathbf{k}_2|^2/2}]^{\frac{1}{2}}}, \quad (25)$$

where $\Delta\mathbf{k}_i \equiv \mathbf{k}_L - k_L \mathbf{r}_i / |\mathbf{r}_i|$ ($i = 1$ and 2). When the number of atoms is sufficiently large to satisfy $N e^{-|d\Delta\mathbf{k}_i|^2/2} \gg 1$ for $i = 1$ and 2 , the degree of the first-order coherence becomes $|g^{(1)}(\mathbf{r}_1, t_1; \mathbf{r}_2, t_2)| \simeq 1$. The scattered photon field is therefore first-order coherent when the number of atoms in the condensate is sufficiently large. On the other hand, when $N e^{-|d\Delta\mathbf{k}_i|^2/2} \ll 1$ for both $i = 1$ and 2 , Eq. (25) reduces to

$$|g^{(1)}(\mathbf{r}_1, t_1; \mathbf{r}_2, t_2)| \simeq \exp\left(-\frac{d^2}{4} |\Delta\mathbf{k}_1 e^{-i\omega(t_1-t_2)} - \Delta\mathbf{k}_2|^2\right). \quad (26)$$

When $t_1 = t_2$, the coherence (26) is lost exponentially with increasing the difference in the scattering angles $|\Delta\mathbf{k}_1 - \Delta\mathbf{k}_2|$. This is because two photon paths corresponding to atoms kicked in different directions do not interfere with each other. The coherence is periodic with respect to the time interval $t_1 - t_2$ with the period of $2\pi\omega^{-1}$. We note that the same expression as Eq. (26) can be obtained by substituting $N = 1$ in Eq. (25), and therefore the correlation described above cannot be ascribed to BEC.

Figure 2 shows the degree of first-order coherence for $t_1 = t_2$, $|g^{(1)}(\mathbf{r}_1, t; \mathbf{r}_2, t)|$, where $N = 10^3$, $d = 1\mu\text{m}$, and $2\pi/k_L = 589\text{nm}$ (the 3s-3p transition of ^{23}Na) are




FIG. 2. The angular dependence of the degree of first-order coherence $g^{(1)}(\mathbf{r}_1, t; \mathbf{r}_2, t)$ of photons scattered from 10^3 BEC atoms. The wavelength of the laser field $2\pi/k_L = 589\text{nm}$ (the 3s-3p transition of ^{23}Na) and the characteristic length of the trap $d = 1\mu\text{m}$ are assumed. The wave vectors $\mathbf{k}_1$ and $\mathbf{k}_2$ are assumed to be in the same plane, and θ_1 and θ_2 denote their angles as measured from $\mathbf{k}_L$.

assumed. The wave vector $\mathbf{k}_1$ and $\mathbf{k}_2$ are assumed to be in the same plane, and θ_1 and θ_2 denote the angles of $\mathbf{k}_1$ and $\mathbf{k}_2$ as measured from $\mathbf{k}_L$ (see Fig. 1). One can see that $g^{(1)} \simeq 1$ for $|d\Delta\mathbf{k}| \lesssim 1$, and therefore, the photons scattered in the forward directions have first-order coherence. There are peaks along $\theta_1 = \theta_2$, which can be seen from Eq. (26).

The degree of second-order coherence [24] is defined by

$$g^{(2)}(\mathbf{r}_1, t_1; \mathbf{r}_2, t_2) \equiv \frac{\sum_{\alpha, \beta=x, y, z} \langle \hat{E}_\alpha^{(-)}(\mathbf{r}_1, t_1) \hat{E}_\beta^{(-)}(\mathbf{r}_2, t_2) \hat{E}_\beta^{(+)}(\mathbf{r}_2, t_2) \hat{E}_\alpha^{(+)}(\mathbf{r}_1, t_1) \rangle}{I(\mathbf{r}_1, t_1) I(\mathbf{r}_2, t_2)}. \quad (27)$$

Substituting Eq. (17) into this yields

$$g^{(2)}(\mathbf{r}_1, t_1; \mathbf{r}_2, t_2) = \left\{ N(N-1)(N-2)(N-3) e^{-|d\Delta\mathbf{k}_1|^2/2 - |d\Delta\mathbf{k}_2|^2/2} + N(N-1)(N-2) \left[2e^{-|d\Delta\mathbf{k}_1|^2/2 - |d\Delta\mathbf{k}_2|^2/2} \times \text{Re}\left(e^{-d^2\Delta\mathbf{k}_1 \cdot \Delta\mathbf{k}_2 e^{i\omega(t_2-t_1)/2} + e^{d^2\Delta\mathbf{k}_1 \cdot \Delta\mathbf{k}_2 e^{-i\omega(t_2-t_1)/2}}\right) + e^{-|d\Delta\mathbf{k}_1|^2/2} + e^{-|d\Delta\mathbf{k}_2|^2/2} \right] + N(N-1) \left[1 + e^{-d^2|\Delta\mathbf{k}_1/2 - \Delta\mathbf{k}_2 e^{i\omega(t_2-t_1)}|^2} + e^{-d^2|\Delta\mathbf{k}_1 + \Delta\mathbf{k}_2 e^{i\omega(t_2-t_1)}|^2/2} + 2e^{-|d\Delta\mathbf{k}_1|^2/2} + 2e^{-|d\Delta\mathbf{k}_2|^2/2} \cos(d^2\Delta\mathbf{k}_1 \cdot \Delta\mathbf{k}_2 \sin \omega(t_2 - t_1)) \right] + N \right\} / \left\{ N^2 \left[1 + (N-1)e^{-|d\Delta\mathbf{k}_1|^2/2} \right] \left[1 + (N-1)e^{-|d\Delta\mathbf{k}_2|^2/2} \right] \right\}. \quad (28)$$

If the number of atoms is sufficiently large to satisfy $Ne^{-|d\Delta\mathbf{k}_i|^2/2} \gg 1$ for both $i = 1$ and 2 , the degree of second-order coherence becomes $g^{(2)}(\mathbf{r}_1, t_1; \mathbf{r}_2, t_2) \simeq 1$ as well as $|g^{(1)}(\mathbf{r}_1, t_1; \mathbf{r}_2, t_2)| \simeq 1$. In general, r th-order coherence [24] becomes unity for all r , and therefore, in this large N limit the scattered photons are coherent in all orders. On the other hand, if $Ne^{-|d\Delta\mathbf{k}_i|^2/2} \ll 1$ for both $i = 1$ and 2 , Eq. (28) reduces to

$$g^{(2)}(\mathbf{r}_1, t_1; \mathbf{r}_2, t_2) \simeq 1 + e^{-d^2|\Delta\mathbf{k}_1 - \Delta\mathbf{k}_2 e^{i\omega(t_2 - t_1)}|^2/2} + e^{-d^2|\Delta\mathbf{k}_1 + \Delta\mathbf{k}_2 e^{i\omega(t_2 - t_1)}|^2/2}. \quad (29)$$

When $t_1 = t_2$, Eq. (29) becomes $g^{(2)} \simeq 2$ for either $\Delta\mathbf{k}_1 \simeq \Delta\mathbf{k}_2$ or $\Delta\mathbf{k}_1 \simeq -\Delta\mathbf{k}_2$. This means that a pair of photons tends to be scattered either in the same direction or in the symmetrical directions with respect to laser propagation. These correlation in scattered photons can be interpreted as follows. In the case of $\Delta\mathbf{k}_1 \simeq \Delta\mathbf{k}_2$, the first photon kicks an atom from the condensate to the state $e^{i\Delta\mathbf{k}_1 \cdot \mathbf{R}} f_0(\mathbf{R})$, and so does the subsequent photon, where $f_0(\mathbf{R})$ is the ground state of the trap potential. Consequently, the Bose-enhancement factor of this process is

$$\langle (N-2)_0, 2_{\Delta\mathbf{k}_1} | \hat{b}_{\Delta\mathbf{k}_2}^\dagger \hat{b}_0^\dagger \hat{b}_{\Delta\mathbf{k}_1}^\dagger \hat{b}_0 | N_0, 0_{\Delta\mathbf{k}_1} \rangle \simeq \sqrt{2N(N-1)}, \quad (30)$$

where $\hat{b}_{\mathbf{k}}^\dagger \equiv \sum_n \langle n | e^{i\mathbf{k} \cdot \mathbf{R}} | 0 \rangle \hat{b}_n^\dagger$ and $|m_{\mathbf{k}}\rangle \equiv \hat{b}_{\mathbf{k}}^{\dagger m} / \sqrt{m!} |0\rangle$. Therefore the probability that this process occurs is enhanced by a factor of two. In the case of $\Delta\mathbf{k}_1 \simeq -\Delta\mathbf{k}_2$, there is a process in which the subsequent photon kicks back the atom excited by the first photon to the condensate, in addition to the process in which two atoms are excited to the states $e^{i\Delta\mathbf{k}_1 \cdot \mathbf{R}} f_0(\mathbf{R})$ and $e^{i\Delta\mathbf{k}_2 \cdot \mathbf{R}} f_0(\mathbf{R})$. The Bose-enhancement factor of each process is

$$\langle N_0, 0_{\Delta\mathbf{k}_1} | \hat{b}_0^\dagger \hat{b}_{\Delta\mathbf{k}_1}^\dagger \hat{b}_{\Delta\mathbf{k}_1}^\dagger \hat{b}_0 | N_0, 0_{\Delta\mathbf{k}_1} \rangle \simeq N \quad (31)$$

and

$$\langle (N-2)_0, 1_{\Delta\mathbf{k}_1}, 1_{\Delta\mathbf{k}_2} | \hat{b}_0^\dagger \hat{b}_{\Delta\mathbf{k}_1}^\dagger \hat{b}_{\Delta\mathbf{k}_1}^\dagger \hat{b}_0 | N_0, 0_{\Delta\mathbf{k}_1}, 0_{\Delta\mathbf{k}_2} \rangle \simeq \sqrt{N(N-1)}, \quad (32)$$

and hence the probability is enhanced by a factor of two for large N . Thus the above correlations are due to the quantum-statistical effect of bosons. In fact, the degree of second-order coherence reduces to $g^{(2)} = 1$ when $N = 1$, and a photon pair scattered from a single atom shows no correlation. We note that the presence of the condensate is required for the above discussion. The process in which the condensate participates is accompanied by the Bose-enhancement factor of $\sqrt{N}$, and the probability that the process occurs is N times greater than that of the process between other levels. Therefore, scattering of photons by the condensate can clearly be distinguished from that by the non-condensed atoms. When the condensate is not




FIG. 3. The angular dependence of the degree of second-order coherence $g^{(2)}(\mathbf{r}_1, t; \mathbf{r}_2, t)$ of photons scattered from 10^3 BEC atoms. The wavelength of the laser field $2\pi/k_L = 589\text{nm}$ (the 3s-3p transition of ^{23}Na) and the characteristic length of the trap $d = 1\mu\text{m}$ are assumed. The wave vectors $\mathbf{k}_1$ and $\mathbf{k}_2$ are assumed to be in the same plane, and θ_1 and θ_2 denote their angles as measured from $\mathbf{k}_L$.

present and the atoms are thermally distributed, occupation numbers of any states are smaller than one, and Bose-enhancement effects cannot be observed.

The degree of second-order coherence (28) for $t_1 = t_2$ is shown in Fig. 3, where $N = 10^3$ and $d = 1\mu\text{m}$, and $\mathbf{k}_1$ and $\mathbf{k}_2$ are assumed to be on the same plane. We find that $g^{(2)} \simeq 1$ when the scattering angles are small, and $g^{(2)}$ grows with increasing θ_1 . While the peak of $g^{(2)}$ at the same scattering angle $\theta_1 = \theta_2$ is $g^{(2)} \simeq 2$ for large θ_1 , the peak at the opposite angle $\theta_1 = -\theta_2$ first grows and then decreases to $g^{(2)} \simeq 1$ as θ_1 increases. This is due to the fact that $\Delta\mathbf{k}_1 = -\Delta\mathbf{k}_2$ is not compatible with the energy conservation $|\mathbf{k}_1| = |\mathbf{k}_2| = \omega_L/c$ (the recoil energy is much smaller than the photon energy), and the third term in Eq. (29) cannot become unity even for $\theta_1 = -\theta_2$. In other words, the energy conservation prohibits the second photon from exactly kicking the atom back to the condensate.

Figure 4 shows the time dependence of the degree of second-order coherence (28). The parameters are the same as those in Fig. 3, and the scattering angle of the first photon is fixed at $\theta_1 = \pi/12$. We find that the second-order coherence is periodic in $t_2 - t_1$ with the period given by $\pi\omega^{-1}$. This periodic behavior reflects the fact that a recoil atom oscillates in the trap. The coherence disappears when the atom is kicked out of the condensate, and revives when the atom returns to the condensate. We note that $g^{(2)}$ goes to below unity before and after the appearance of the peaks. This indicates that photons are suppressed to be scattered into $\theta_2 \simeq \pm\theta_1$ in these time intervals. This comes from the terms proportional to $N(N-1)(N-2)$ in the numera-




FIG. 4. The time and angular dependence of the degree of second-order coherence $g^{(2)}(\mathbf{r}_1, t_1; \mathbf{r}_2, t_2)$ of photons scattered from 10^3 BEC atoms. The scattering angle of the first photon is fixed at $\theta_1 = \pi/12$. The wavelength of the laser field $2\pi/k_L = 589\text{nm}$ (the 3s-3p transition of ^{23}Na) and the characteristic length of the trap $d = 1\mu\text{m}$ are assumed. The wave vectors $\mathbf{k}_1$ and $\mathbf{k}_2$ are assumed to be in the same plane, and θ_1 and θ_2 denote their angles as measured from $\mathbf{k}_L$.

tor of Eq. (28), which shows destructive interference of probability amplitudes of two-photon processes in which one photon kicks an atom and another photon does not disturb the atoms.

The degree of first-order coherence $g^{(1)}$ can be measured by superposing the photon field in the directions of $\mathbf{k}_1$ and $\mathbf{k}_2$ with a delay time $t_2 - t_1$ in the path of the $\mathbf{k}_1$ photon. Taking a delay of order $\omega^{-1} \sim 1$ ms while retaining the coherence might be feasible by reducing the speed of light using an ultracold atomic gas [9]. On the other hand, a measurement of $g^{(2)}$ does not require the coherent delay, since it measures a correlation between detected photons.

IV. EFFECTS OF ATOM-ATOM INTERACTIONS

In this section, we take into account the atom-atom interaction on the coherence properties of photons scattered from BEC within the Bogoliubov approximation [23].

The second-quantized form of the kinetic part of the Hamiltonian is given by

$$\hat{K} = \int d\mathbf{R} \hat{\psi}^\dagger(\mathbf{R}) \left[-\frac{\hbar^2}{2m} \nabla^2 + V_t(\mathbf{R}) - \mu \right] \hat{\psi}(\mathbf{R}) + \frac{U_0}{2} \int d\mathbf{R} \hat{\psi}^\dagger(\mathbf{R}) \hat{\psi}^\dagger(\mathbf{R}) \hat{\psi}(\mathbf{R}) \hat{\psi}(\mathbf{R}), \quad (33)$$

where $\hat{\psi}(\mathbf{R})$ is the field operator of the atoms (18) and $U_0 \equiv 4\pi\hbar^2 a/m$, a being the s-wave scattering length. In

the Bogoliubov approximation, the atomic field operator is divided into a mean field part and fluctuations from it as

$$\hat{\psi}(\mathbf{R}) = \sqrt{N} \psi_g(\mathbf{R}) + \delta\hat{\psi}(\mathbf{R}), \quad (34)$$

where $\psi_g(\mathbf{R})$ is the normalized single-particle wave function which obeys the Gross-Pitaevskii (GP) equation [25]

$$\left[-\frac{\hbar^2}{2m} \nabla^2 + V_t(\mathbf{R}) + U_0 N |\psi(\mathbf{R})|^2 \right] \psi(\mathbf{R}) = \mu \psi(\mathbf{R}), \quad (35)$$

where μ is determined so that the solution of Eq. (35) satisfies the normalization condition $\int d\mathbf{R} |\psi(\mathbf{R})|^2 = 1$. Equation (35) guarantees that if we substitute Eq. (34) into Eq. (33), terms linear in $\delta\hat{\psi}$ or $\delta\hat{\psi}^\dagger$ in Eq. (33) vanish identically. Because the interaction between atoms is weak, we may expect only small deviations from the mean field at zero temperature, so that we may keep only terms up to quadratic in $\delta\hat{\psi}$ and $\delta\hat{\psi}^\dagger$:

$$\begin{aligned} \hat{K} \simeq & N \int d\mathbf{R} \psi_g^*(\mathbf{R}) \left[-\frac{\hbar^2}{2m} \nabla^2 + V_t(\mathbf{R}) - \mu \right. \\ & \left. + \frac{U_0 N}{2} |\psi(\mathbf{R})|^2 \right] \psi_g(\mathbf{R}) \\ & + \int d\mathbf{R} \left[\delta\hat{\psi}^\dagger(\mathbf{R}) \left(-\frac{\hbar^2}{2m} \nabla^2 + V_t(\mathbf{R}) - \mu \right. \right. \\ & \left. \left. + 2U_0 N |\psi(\mathbf{R})|^2 \right) \delta\hat{\psi}(\mathbf{R}) \right. \\ & \left. + U_0 N \psi_g^{*2}(\mathbf{R}) \delta\hat{\psi}^2(\mathbf{R}) + U_0 N \psi_g^2(\mathbf{R}) \delta\hat{\psi}^{\dagger 2}(\mathbf{R}) \right] \quad (36) \end{aligned}$$

To diagonalize the quadratic terms, we write $\delta\hat{\psi}(\mathbf{R})$ as

$$\delta\hat{\psi}(\mathbf{R}) = \sum_{\lambda} \left[u_{\lambda}(\mathbf{R}) \hat{\beta}_{\lambda} + v_{\lambda}^*(\mathbf{R}) \hat{\beta}_{\lambda}^{\dagger} \right]. \quad (37)$$

Requiring the coefficients of the terms $\hat{\beta}_{\lambda}^2$ and $\hat{\beta}_{\lambda}^{\dagger 2}$ to vanish, we find that $u_{\lambda}(\mathbf{R})$ and $v_{\lambda}(\mathbf{R})$ should satisfy

$$\left[-\frac{\hbar^2}{2m} \nabla^2 + V_t(\mathbf{R}) - \mu + 2U_0 N |\psi_g(\mathbf{R})|^2 \right] u_{\lambda}(\mathbf{R}) + U_0 N \psi_g(\mathbf{R})^2 v_{\lambda}(\mathbf{R}) = \hbar\omega_{\lambda} u_{\lambda}(\mathbf{R}), \quad (38a)$$

$$\left[-\frac{\hbar^2}{2m} \nabla^2 + V_t(\mathbf{R}) - \mu + 2U_0 N |\psi_g(\mathbf{R})|^2 \right] v_{\lambda}(\mathbf{R}) + U_0 N \psi_g^{*2}(\mathbf{R}) u_{\lambda}(\mathbf{R}) = -\hbar\omega_{\lambda} v_{\lambda}(\mathbf{R}). \quad (38b)$$

The quadratic part of the Hamiltonian (36) can then be diagonalized as

$$\hat{K} = \sum_{\lambda} \hbar\omega_{\lambda} \hat{\beta}_{\lambda}^{\dagger} \hat{\beta}_{\lambda}, \quad (39)$$

where $\hat{\beta}_{\lambda}^{\dagger}$, $\hat{\beta}_{\lambda}$ are the creation and annihilation operators of the Bogoliubov quasiparticles and $\hbar\omega_{\lambda}$ denotes the energy of elementary excitations. The constant terms are

irrelevant to later discussions and are therefore omitted in Eq. (39).

The second quantized form of the operator $\sum_{j=1}^N e^{i\Delta\mathbf{k}\cdot\mathbf{R}_j(t)}$ that gives the momentum to an atom reads in the Bogoliubov approximation as

$$\hat{\mathcal{D}}_B(\Delta\mathbf{k}, t) \equiv \int d\mathbf{R} \hat{\psi}^\dagger(\mathbf{R}, t) e^{i\Delta\mathbf{k}\cdot\mathbf{R}} \hat{\psi}(\mathbf{R}, t), \quad (40)$$

where $\hat{\psi}(\mathbf{R}, t) = e^{\frac{i}{\hbar}\hat{K}t}\hat{\psi}(\mathbf{R})e^{-\frac{i}{\hbar}\hat{K}t}$. Because $|u_\lambda(\mathbf{R})| \gg |v_\lambda(\mathbf{R})|$ [26], we may neglect $v_\lambda(\mathbf{R})$ in Eq. (34). The operator (40) then becomes

$$\begin{aligned} \hat{\mathcal{D}}_B(\Delta\mathbf{k}, t) = & N\langle\psi_g|e^{i\Delta\mathbf{k}\cdot\mathbf{R}}|\psi_g\rangle \\ & + \sqrt{N} \sum_{\lambda} \left(\langle\psi_g|e^{i\Delta\mathbf{k}\cdot\mathbf{R}}|u_\lambda\rangle \hat{\beta}_\lambda e^{-i\omega_\lambda t} \right. \\ & \left. + \langle u_\lambda|e^{i\Delta\mathbf{k}\cdot\mathbf{R}}|\psi_g\rangle \hat{\beta}_\lambda^\dagger e^{i\omega_\lambda t} \right) \\ & + \sum_{\lambda\mu} \langle u_\lambda|e^{i\Delta\mathbf{k}\cdot\mathbf{R}}|u_\mu\rangle \hat{\beta}_\lambda^\dagger \hat{\beta}_\mu e^{i(\omega_\lambda - \omega_\mu)t}, \quad (41) \end{aligned}$$

where we use the shorthand notation as in Eq. (20). In the Bogoliubov approximation, the orthonormality condition is expressed as $\langle u_\lambda|u_\mu\rangle - \langle v_\lambda|v_\mu\rangle = \delta_{\lambda\mu}$, and the completeness relation is given by $\sum_{\lambda} (|u_\lambda\rangle\langle u_\lambda| - |v_\lambda\rangle\langle v_\lambda|) = 1 - |\psi_g\rangle\langle\psi_g|$. In the present approximation of neglecting v_λ , they reduce to $\langle u_\lambda|u_\mu\rangle = \delta_{\lambda\mu}$ and $\sum_{\lambda} |u_\lambda\rangle\langle u_\lambda| + |\psi_g\rangle\langle\psi_g| = 1$, respectively.

Using the operator (41), the source-field operator is expressed as $\hat{\mathbf{E}}_s^{(+)}(\mathbf{r}, t) = \mathbf{F}(\mathbf{r}, t)\hat{\mathcal{D}}_B(\Delta\mathbf{k}, t)$. The expectation values of the electric field and the intensity are calculated to be

$$\langle \hat{\mathbf{E}}^{(+)}(\mathbf{r}, t) \rangle = \mathbf{F}(\mathbf{r}, t) N \langle \psi_g | e^{i\Delta\mathbf{k}\cdot\mathbf{R}} | \psi_g \rangle, \quad (42)$$

$$I(\mathbf{r}, t) = |\mathbf{F}(\mathbf{r}, t)|^2 N (1 + N |\langle \psi_g | e^{i\Delta\mathbf{k}\cdot\mathbf{R}} | \psi_g \rangle|^2). \quad (43)$$

These expressions are different from the noninteracting counterparts (21) and (23) in that the expectation value with respect to the ground state of the noninteracting atoms $\langle 0 | e^{i\Delta\mathbf{k}\cdot\mathbf{R}} | 0 \rangle$ is replaced by that of the ground state of the GP equation $\langle \psi_g | e^{i\Delta\mathbf{k}\cdot\mathbf{R}} | \psi_g \rangle$, and $N - 1$ by N . The latter arises from the assumption of the Bogoliubov theory that the ground state is a coherent state rather than a Fock state. Although this is an artifact, the error is of order $1/N$ and negligible for $N \gg 1$. The former reflects the fact that the ground-state wave function of interacting atoms expands for the repulsive interaction $a > 0$, and contracts for the attractive interaction $a < 0$, compared with that of noninteracting atoms.

The degree of first-order coherence is calculated to be

$$\begin{aligned} |g^{(1)}(\mathbf{r}_1, t_1; \mathbf{r}_2, t_2)| = & \frac{\langle e^{-i\Delta\mathbf{k}_1\cdot\mathbf{R}} \hat{U}_B(t_1 - t_2) e^{i\Delta\mathbf{k}_2\cdot\mathbf{R}} \rangle_g + N \langle e^{-i\Delta\mathbf{k}_1\cdot\mathbf{R}} \rangle_g \langle e^{i\Delta\mathbf{k}_2\cdot\mathbf{R}} \rangle_g}{(1 + N |\langle e^{i\Delta\mathbf{k}_1\cdot\mathbf{R}} \rangle_g|^2)^{\frac{1}{2}} (1 + N |\langle e^{i\Delta\mathbf{k}_2\cdot\mathbf{R}} \rangle_g|^2)^{\frac{1}{2}}}, \quad (44) \end{aligned}$$




FIG. 5. The angular dependence of the degree of first-order coherence $g^{(1)}(\mathbf{r}_1, t; \mathbf{r}_2, t)$ of photons scattered from 10^3 interacting BEC atoms. The s-wave scattering length is assumed to be $a = 2.75$ nm. The wavelength of the laser field $2\pi/k_L = 589$ nm (the 3s-3p transition of ^{23}Na) and the characteristic length of the trap $d = 1\mu\text{m}$ are assumed. The wave vectors $\mathbf{k}_1$ and $\mathbf{k}_2$ are assumed to be in the same plane, and θ_1 and θ_2 denote their angles as measured from $\mathbf{k}_L$.

where $\hat{U}_B(t) \equiv \sum_{\lambda} |u_\lambda\rangle e^{-i\omega_\lambda t} \langle u_\lambda|$, and $\langle \cdots \rangle_g$ denotes the expectation value with respect to the ground state of the GP equation (35). To evaluate the expectation value that includes $\hat{U}_B$, we need to find u_λ which we obtain by numerically diagonalizing Eqs. (38a) and (38b) using the method discussed in Ref. [27]. Figure 5 shows the degree of first-order coherence (44) for $t_1 = t_2$. The parameters are taken to be $N = 10^3$, $d = 1\mu\text{m}$, and $a = 2.75$ nm. The ground state of the GP equation (35) is obtained numerically [22]. The sharp dips in Fig. 5 correspond to the points at which the sign of $g^{(1)}$ changes (note that Fig. 5 displays the absolute value of $g^{(1)}$), while the curves in the noninteracting case are smooth (see Fig. 2). To understand the oscillatory behaviors of $g^{(1)}$, we compare in Fig. 6 the profile of the interacting ground-state wave function with $N = 10^3$ with the corresponding noninteracting one. As we can see from the figure the density profile near the origin becomes flat due to the repulsive interaction which gives rise to the oscillatory behavior of $e^{i\Delta\mathbf{k}\cdot\mathbf{R}}$. The validity of this interpretation can be confirmed analytically in the limit of strong interaction $Na/d \gg 1$. In this limit the kinetic energy may be neglected in comparison with the trap potential and the interaction energy (Thomas-Fermi approximation), and the wave function is given by

$$\psi_{\text{TF}}(R) = \sqrt{\frac{15}{8\pi R_0^5}} (R_0^2 - R^2), \quad (45)$$

where $R_0 \equiv d(15Na/d)^{1/5}$ [28]. In this Thomas-Fermi limit, the expectation value becomes




FIG. 6. The profiles of the ground-state wave functions. The solid curve shows the wave function of interacting 10^3 atoms which is obtained by solving the GP equation (35), and the dashed curve shows the noninteracting one.

$$\langle \psi_g | e^{i\Delta \mathbf{k} \cdot \mathbf{R}} | \psi_g \rangle = \frac{15}{2|R_0 \Delta \mathbf{k}|^5} [(6 - 2R_0 |\Delta \mathbf{k}|) \sin R_0 |\Delta \mathbf{k}| - 6R_0 |\Delta \mathbf{k}| \cos R_0 |\Delta \mathbf{k}|], \quad (46)$$

which shows oscillations of $g^{(1)}$. In contrast the Gaussian wave function gives a monotonically decreasing nonzero value $\langle e^{i\Delta \mathbf{k} \cdot \mathbf{R}} \rangle = e^{-|d\Delta \mathbf{k}|^2/4}$.

The degree of second-order coherence is calculated to be

$$\begin{aligned} g^{(2)}(\mathbf{r}_1, t_1; \mathbf{r}_2, t_2) = & \left[N^4 |\langle e^{i\Delta \mathbf{k}_1 \cdot \mathbf{R}} \rangle_g \langle e^{i\Delta \mathbf{k}_2 \cdot \mathbf{R}} \rangle_g|^2 \right. \\ & + N^3 \left(2\text{Re} \langle e^{-i\Delta \mathbf{k}_1 \cdot \mathbf{R}} \rangle_g \langle e^{-i\Delta \mathbf{k}_2 \cdot \mathbf{R}} \rangle_g \right. \\ & \times \langle e^{i\Delta \mathbf{k}_2 \cdot \mathbf{R}} \hat{U}_B(t_2 - t_1) e^{i\Delta \mathbf{k}_1 \cdot \mathbf{R}} \rangle_g \\ & + 2\text{Re} \langle e^{-i\Delta \mathbf{k}_1 \cdot \mathbf{R}} \rangle_g \langle e^{i\Delta \mathbf{k}_2 \cdot \mathbf{R}} \rangle_g \langle e^{-i\Delta \mathbf{k}_2 \cdot \mathbf{R}} \hat{U}_B(t_2 - t_1) e^{i\Delta \mathbf{k}_1 \cdot \mathbf{R}} \rangle_g \\ & \left. \left. + |\langle e^{i\Delta \mathbf{k}_1 \cdot \mathbf{R}} \rangle_g|^2 + |\langle e^{i\Delta \mathbf{k}_2 \cdot \mathbf{R}} \rangle_g|^2 \right) \right. \\ & + N^2 \left(1 + |\langle e^{i\Delta \mathbf{k}_2 \cdot \mathbf{R}} \hat{U}_B(t_2 - t_1) e^{i\Delta \mathbf{k}_1 \cdot \mathbf{R}} \rangle_g|^2 \right. \\ & + |\langle e^{-i\Delta \mathbf{k}_2 \cdot \mathbf{R}} \hat{U}_B(t_2 - t_1) e^{i\Delta \mathbf{k}_1 \cdot \mathbf{R}} \rangle_g|^2 \\ & + 2|\langle e^{i\Delta \mathbf{k}_1 \cdot \mathbf{R}} \rangle_g|^2 + 2\text{Re} \langle e^{-i\Delta \mathbf{k}_2 \cdot \mathbf{R}} \rangle_g \\ & \times \langle e^{-i\Delta \mathbf{k}_1 \cdot \mathbf{R}} \hat{U}_B^\dagger(t_2 - t_1) e^{i\Delta \mathbf{k}_2 \cdot \mathbf{R}} \hat{U}_B(t_2 - t_1) e^{i\Delta \mathbf{k}_1 \cdot \mathbf{R}} \rangle_g \\ & \left. \left. + N \right] \right] / \left[N^2 (1 + N |\langle e^{i\Delta \mathbf{k}_1 \cdot \mathbf{R}} \rangle_g|^2) (1 + N |\langle e^{i\Delta \mathbf{k}_2 \cdot \mathbf{R}} \rangle_g|^2) \right]. \end{aligned} \quad (47)$$

Figure 7 shows the degree of second-order coherence (47) for $t_1 = t_2$, where the parameters are the same as in Fig. 5. One can see the complicated structure that arises




FIG. 7. The angular dependence of the degree of second-order coherence $g^{(2)}(\mathbf{r}_1, t; \mathbf{r}_2, t)$ of photons scattered from 10^3 interacting BEC atoms. The s-wave scattering length is assumed to be $a = 2.75$ nm. The wavelength of a laser field $2\pi/k_L = 589$ nm (the 3s-3p transition of ^{23}Na) and the characteristic length of the trap $d = 1 \mu\text{m}$ are assumed. The wave vectors $\mathbf{k}_1$ and $\mathbf{k}_2$ are assumed to be in the same plane, and θ_1 and θ_2 denote their angles as measured from $\mathbf{k}_L$.

from the oscillations of $\langle e^{i\Delta \mathbf{k}_1 \cdot \mathbf{R}} \rangle_g$. Figure 8 shows the dependence on the time interval $t_2 - t_1$ of $g^{(2)}(\mathbf{r}_1, t_1; \mathbf{r}_2, t_2)$, where the parameters are the same as in Fig. 5, and the scattering angle of the first photon θ_1 is fixed at $\pi/12$. We find that the peaks appear periodically with their heights monotonically decreasing. The decay of the peaks can be understood as follows. The atom is kicked out of the condensate upon receipt of recoil momentum, and oscillates in the trap potential while encountering the condensate periodically. If the atom-atom interaction is not present, the Gaussian wave function of the scattered atom retains its shape during the oscillations, and the peaks revive completely as shown in Fig. 4. On the other hand, when the atom-atom interaction is present, the wave function undergoes distortion by colliding the condensate. As a result, the atom cannot be kicked back completely to the condensate by scattering a photon, and therefore the correlation deteriorates.

V. CONCLUSIONS

We have studied the coherence properties of photons scattered from the BEC irradiated by a weak and far-off-resonant laser field, and have shown that the degree of second-order coherence $g^{(2)}$ exhibits unique features of trapped BEC. That is, the probability of two photons being scattered either in the same direction or in the symmetrical direction with respect to laser propagation is enhanced, and these correlations appear periodically with the period of atomic motion in the trapping potential.




FIG. 8. The time and angular dependence of the degree of second-order coherence $g^{(2)}(\mathbf{r}_1, t_1; \mathbf{r}_2, t_2)$ of photons scattered from 10^3 interacting BEC atoms. The scattering angle of the first photon is fixed at $\theta_1 = \pi/12$. The s-wave scattering length is assumed to be $a = 2.75$ nm. The wavelength of the laser field $2\pi/k_L = 589$ nm (the 3s-3p transition of ^{23}Na) and the characteristic length of the trap $d = 1\mu\text{m}$ are assumed. The wave vectors $\mathbf{k}_1$ and $\mathbf{k}_2$ are assumed to be in the same plane, and θ_1 and θ_2 denote their angles as measured from $\mathbf{k}_L$.

These correlations in space and time are manifestations of the combined effect of bosonic stimulation and quantized kinetic motion in the trap. We have also taken into account effects of the atom-atom interaction, and have confirmed that such correlations can be observed even in the presence of the atom-atom interaction with appearance of an additional structure (sharp dips in Fig. 5) resulting from the repulsive nature of the interaction.

ACKNOWLEDGMENT

One of the authors (H.S.) acknowledges support by the Japan Society for the Promotion of Science for Young Scientists.

-
- [1] M. H. Anderson, J. R. Ensher, M. R. Matthews, C. E. Wieman, and E. A. Cornell, *Science* **269**, 198 (1995).
 - [2] C. C. Bradley, C. A. Sackett, J. J. Tollett, and R. G. Hulet, *Phys. Rev. Lett.* **75**, 1687 (1995).
 - [3] K. B. Davis, M. -O. Mewes, M. R. Andrews, N. J. van Druten, D. S. Durfee, D. M. Kurn, and W. Ketterle, *Phys. Rev. Lett.* **75**, 3969 (1995).
 - [4] D. G. Fried, T. C. Killian, L. Willmann, D. Landhuis,

- S. C. Moss, D. Kleppner, and T. J. Greytak, *Phys. Rev. Lett.* **81**, 3811 (1998).
- [5] M. R. Andrews, M. -O. Mewes, N. J. van Druten, D. S. Durfee, D. M. Kurn, and W. Ketterle, *Science* **273**, 84 (1996).
- [6] M. R. Andrews, C. G. Townsend, H. -J. Miesner, D. S. Durfee, D. M. Kurn, and W. Ketterle, *Science* **275**, 637 (1997).
- [7] M. R. Andrews, D. M. Kurn, H. -J. Miesner, D. S. Durfee, C. G. Townsend, S. Inouye, and W. Ketterle, *Phys. Rev. Lett.* **79**, 553 (1997).
- [8] D. S. Hall, M. R. Matthews, J. R. Ensher, C. E. Wieman, and E. A. Cornell, *Phys. Rev. Lett.* **81**, 1539 (1998).
- [9] L. V. Hau, S. E. Harris, Z. Dutton, and C. H. Behroozi, *Nature* **397**, 594 (1999).
- [10] B. V. Svistunov and G. V. Shlyapnikov, *Zh. Eksp. Teor. Fiz.* **97**, 821 (1990); **98**, 129 (1990) [*Sov. Phys. JETP* **70**, 460 (1990); **71**, 71 (1990)].
- [11] H. D. Politzer, *Phys. Rev. A* **43**, 6444 (1991).
- [12] M. Lewenstein and L. You, *Phys. Rev. Lett.* **71**, 1339 (1993).
- [13] J. Javanainen, *Phys. Rev. Lett.* **72**, 2375 (1994).
- [14] M. Lewenstein, L. You, J. Cooper, and K. Burnett, *Phys. Rev. A* **50**, 2207, (1994).
- [15] L. You, M. Lewenstein, and J. Cooper, *Phys. Rev. A* **50**, R3565 (1994).
- [16] J. Javanainen, *Phys. Rev. Lett.* **75**, 1927 (1995),
- [17] L. You, M. Lewenstein, and J. Cooper, *Phys. Rev. A* **51**, 4712 (1995).
- [18] O. Morice, Y. Castin, and J. Dalibard, *Phys. Rev. A* **51**, 3896 (1995).
- [19] J. Javanainen and J. Ruostekoski, *Phys. Rev. A* **52**, 3033 (1995).
- [20] L. You, M. Lewenstein, R. J. Glauber, and J. Cooper, *Phys. Rev. A* **53**, 329 (1996).
- [21] P. A. Ruprecht, M. J. Holland, K. Burnett, and M. Edwards, *Phys. Rev. A* **51**, 4704 (1995).
- [22] F. Dalfovo and S. Stringari, *Phys. Rev. A* **53**, 2477 (1996).
- [23] N. Bogoliubov, *J. Phys. (USSR)* **11**, 23 (1947).
- [24] R. J. Glauber, *Quantum Optics and Electronics*, edited by C. DeWitt, A. Blandin, and C. Cohen-Tannoudji, (Gordon and Breach, New York, 1965); see also R. Loudon, *The Quantum Theory of Light*, (Oxford, New York, 1983).
- [25] E. P. Gross, *Nuovo Cimento* **20**, 454 (1961); *J. Math. Phys.* **4**, 195 (1963); L. P. Pitaevskii, *Zh. Eksp. Teor. Fiz.* **40**, 646 (1961) [*Sov. Phys. JETP* **13**, 451 (1961)].
- [26] F. Dalfovo, S. Giorgini, M. Guilleumas, L. Pitaevskii, and S. Stringari, *Phys. Rev. A* **56**, 3840 (1997).
- [27] M. Edwards, R. J. Dodd, C. W. Clark, and K. Burnett, *J. Res. Natl. Inst. Stand. Technol.* **101**, 553 (1996).
- [28] G. Baym and C. J. Pethick, *Phys. Rev. Lett.* **76**, 6 (1996).